



CASE 6

G. Petur Nielsen, M.D.

Department of Pathology

Massachusetts General Hospital

Boston, MA

Professor of Pathology

Harvard Medical School

gnielsen@partners.org








CASE 6


 43 year old presented with pain in the right upper thigh that had recently increased.
 There was a history of minor trauma and injury in the remote past





















Diagnosis

Bizarre parosteal osteochondromatous proliferation (BPOP)






BPOP

- First described by Nora et al.
- 35 patients
- 14-74 years old
- 0.4-3 cm
- Proximal phalanges / metatarsals / metacarpals
- 18 local recurrence 8 locally recurred twice
 - Nora et al. Bizarre parosteal osteochondromatous proliferations of the hands and feet.
 Am J Surg Pathol 1983; 7:245-250









- 65 patients
- 8-73 years old
- 72% involved the small bones
- 27% involved long bones
- 14 with one local recurrence 5 with two 3 with three
 - Menses et al. Bizarre parosteal osteochondromatous proliferations of bone (Nora's lesion). Am J Surg Pathol 1993; 17:691-697





Mass General Hosp



Fibrous tissue

Cartilage

Bone





BPOP

- Initially thought to be a non-neoplastic, reactive process
- t(1;17(q32;q21) has been identified supporting a neoplastic process
- Four cases with inversion of chromosome 7 have been described
 - Nilsson et al et al. Molecular cytogenetic characterization of recurrent translocation breakpoints in bizarre parosteal osteochondromatous proliferation (Nora's lesion). Human Pathology 2004;35:1063-1069
 - Endo et al. Bizarre parosteal osteochondromatous proliferation with a t(1;17) translocation. Virchows Archiv 2005;447:99-102
 - Broehm et al. Bizarre parosteal osteochondromatous proliferation: a new cytogenetic subgroup characterized by inversion of chromosome 7. Cancer Genetics 2013;206:402-405





BPOP – Differential Diagnosis

- Small bones
 - Subungual exostosis
 - Florid reactive periostitis






Subungual Exostosis

- Arises in a subungual location
- t(X;6)(q24-26;q15-21)


Location of Subungual Exostosis Based on 244 Cases



- DaCambra et al. Subungual Exostosis of the Toes: A Systematic Review. Clin Orthop Relat Res 2014;472:1251-1259
- Mertens et al. The t(X;6) in subungual exostosis results in transcriptional deregulation of the gene for insulin receptor substrate 4. Int. J. Cancer 2011;128:487-491












Florid Reactive Periostitis

- Surface lesion of small bones of hands and feet (parosteal fasciitis – unusual myositis ossificans)
- Probably reactive / traumatic process no genetic abnormality has been identified

Spjut et al. Florid reactive periostitis of the tubular bones of the hands and feet. A benign lesion which may simulate osteosarcoma. Am J Surg Pathol 1981;5:423-433












BPOP – Differential Diagnosis

- Long bones
 - Parosteal osteosarcoma
 - Periosteal osteosarcoma
 - Osteochondroma


Parosteal Osteosarcoma



Parosteal Osteosarcoma




Parosteal Osteosarcoma




Periosteal Osteosarcoma





Periosteal Osteosarcoma




Osteochondroma







Osteochondroma





Osteochondroma









CASE 6

- Treated with resection
- No recurrence 11 years later

